

After the Shots...

Acetaminophen and ibuprofen are strong medicines and should ONLY be used when needed. Your child does not need acetaminophen or ibuprofen if s/he is mildly ill. Only give acetaminophen or ibuprofen if your child has a fever (that you checked with a thermometer) and is not eating, sleeping or playing normal. If you give your child acetaminophen or ibuprofen, read the directions on the label. Only use the syringe or cup that comes with the medicine to measure it.

If your child has fever:

- Give your child plenty to drink.
- Keep your child cool. Do not put too many clothes or blankets on your child.
- Give your child a bath in 1-2 inches of lukewarm water.

If your child's leg or arm is swollen, hot, or red:

- Put a clean, cool, wet washcloth on the sore area.
- If redness or pain gets worse after 24 hours, call the doctor.

Call your child's doctor right away if your child:

- Has a temperature of 103° F (39.4°C) or higher
- Is pale or limp
- Has been crying for more than 3 hours
- Has a strange cry (high pitched)
- Is shaking, twitching, or jerking
- Is not active or responding
- If you are worried about how your child looks or feels.




Phone number of your child's doctor _____

How much medicine can I give my child?

Acetaminophen (Tylenol® or another brand) Give every 4-6 hours as needed. Don't give more than five (5) times in 24 hours (unless told to do so by a doctor). Read and follow the directions on the label. Only use the syringe or cup that comes with the medicine to measure it. Ibuprofen dosing information on other side.

6- 11 pounds 0 – 3 months	12- 17 pounds 4 – 11 months	18- 23 pounds 12 - 23 months	24- 35 pounds 2- 3 years	36- 47 pounds 4- 5 years	48 – 59 pounds 6 – 8 years	60 – 71 pounds 9 – 10 years	72 – 95 pounds 11 years
<p>1.25 mL Oral Suspension</p> 	<p>2.5 mL Oral Suspension</p> 	<p>3.75 mL Oral Suspension</p> 	<p>5 mL Oral Suspension</p> 	<p>1 ½ teaspoons Children's Liquid</p> 	<p>2 teaspoons Children's Liquid</p> 	<p>2 ½ teaspoons Children's Liquid</p> 	<p>3 teaspoons Children's Liquid</p> 
<p>WARNING:</p> <ul style="list-style-type: none"> • Never give aspirin to a child (unless you are told to by a doctor). • If you give your child acetaminophen or ibuprofen, do NOT also give cold, allergy or sinus medications with acetaminophen or ibuprofen in them. Your child could get too much medicine. • Do not give adult medicines to children. • Keep medicines out of reach of children. 			<p>OR 3 tablets 80 mg. Children's Tablets</p> 	<p>OR 4 tablets 80 mg. Children's Tablets</p> 	<p>OR 5 tablets 80 mg. Children's Tablets</p> 	<p>OR 6 tablets 80 mg. Children's Tablets</p> 	
			<p>1 teaspoon Children's Liquid</p> 	<p>OR 2 tablets 80 mg. Children's Tablets</p> 	<p>OR 2 ½ tablets 160 mg. Junior Strength Tabs</p> 	<p>OR 2 ½ tablets 160 mg. Junior Strength Tabs</p> 	<p>OR 3 tablets 160 mg. Junior Strength Tabs</p> 

Ibuprofen (Advil®, Motrin® or another brand) may be given every 6-8 hours if needed. Don't give more than four (4) times in 24 hours (unless told to do so by a doctor.) Read and follow the directions on the label. Only use the syringe or cup that comes with the medicine to measure it. Acetaminophen dosing information on other side.

12 – 17 pounds 6 – 11 months	18 – 23 pounds 12 – 23 months	24 – 35 pounds 2 – 3 years	36- 47 pounds 4- 5 years	48- 59 pounds 6- 8 years	60- 71 pounds 9- 10 years	72- 95 pounds 11 years
<p>1.25 mL Infant Drops</p> 	<p>1.875 mL Infant Drops</p> 	<p>1 teaspoon Children's Liquid</p>  <p>OR</p> <p>2 tablets 50 mg. Children's Tablets</p> 	<p>1 ½ teaspoons Children's Liquid</p>  <p>OR</p> <p>3 tablets 50 mg. Children's Tablets</p> 	<p>2 teaspoons Children's Liquid</p>  <p>OR</p> <p>4 tablets 50 mg. Children's Tablets</p>  <p>OR</p> <p>2 tablets 100 mg. Junior Strength Tabs</p> 	<p>2 ½ teaspoons Children's Liquid</p>  <p>OR</p> <p>5 tablets 50 mg. Children's Tablets</p>  <p>OR</p> <p>2 ½ tablets 100 mg. Junior Strength Tabs</p> 	<p>3 teaspoons Children's Liquid</p>  <p>OR</p> <p>6 tablets 50 mg. Children's Tablets</p>  <p>OR</p> <p>3 tablets 100 mg. Junior Strength Tabs</p> 

WARNING:

- **Never give aspirin to a child (unless you are told to by a doctor).**
- **If you give your child acetaminophen or ibuprofen, do NOT also give cold, allergy or sinus medications with acetaminophen or ibuprofen in them. Your child could get too much medicine.**
- **Do not give adult medicines to children.**
- **Keep all medicines out of reach of children.**

